

Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 23 February 2015

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Tuesday, February 17, 2015



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from February 13 to 20: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at: http://www.tpwd.state.tx.us./landwater/water/environconcerns/hab/redtide/status.phtml

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive:

http://tidesandcurrents.noaa.gov/hab/bulletins.html

Conditions Report

Karenia brevis (commonly known as Texas red tide) ranges from not present to very low concentrations along the coast of Texas. No respiratory irritation is expected alongshore Texas Monday, February 23 through Monday, March 2.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

Sampling from Texas A&M University's Imaging FlowCytobot, located on the Port Aransas ship channel, continues to indicate that *Karenia brevis* concentrations range between 'not present' and 'very low a' (TAMU; 2/17-23). For information on area shell-fish restrictions, contact the Texas Department of State Health Services.

Recent MODIS Aqua imagery (2/21, shown left) is obscured by clouds from Sabine Pass to the Rio Grande, preventing analysis. In MODIS Aqua imagery from February 19 (not shown), elevated chlorophyll (1-8 μ g/L) is visible stretching along- and offshore from Sabine Pass to the Rio Grande. Elevated chlorophyll is not necessarily indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of 40 km south from the Port Aransas region from February 21-25.

Keeney, Kavanaugh




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

-2-

Wind Analysis

Port Aransas: North winds (10-25kn, 5-13m/s) today through Tuesday shifting to the northeast (10-20kn, 5-10m/s) Tuesday night. North winds (15-20kn, 8-10m/s) Wednesday. Northeast winds (10-15kn, 5-8m/s) Wednesday night through Thursday night. East winds (15-20kn) Friday.



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).